

**PR-7. CERIC AMMONIUM NITRATE (CAN) CATALYZED SYNTHESIS  
AND  $\alpha$ -GLUCOSIDASE ACTIVITY  
OF SOME NOVEL TETRAHYDROPYRIDINE PHOSPHONATE DERIVATIVES**

N. B. Reddy<sup>1</sup>, G. V. Zyryanov<sup>1,2</sup>


<sup>1</sup>Ural Federal University of the first President of Russia B. N. Yeltsin,  
Mira St., 19, Yekaterinburg, 620002, Russia

<sup>2</sup>I. Ya. Postovsky Institute of Organic Synthesis UB RAS,  
S. Kovalevskoy/Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia

E-mail: drbvreddyn@gmail.com

**Abstract.** A simple and highly efficient multicomponent one-pot synthesis of a series of pharmaceutically interesting functionalized tetrahydropyridine phosphonate derivatives have been developed based on low-cost and environmentally benign cerium ammonium nitrate catalyst via tandem reactions of 1,3-dicarbonyl compounds, diethylphosphoramidate and various aromatic aldehydes in acetonitrile at room temperature. High atom economy, good yields, eco-friendliness and mild reaction conditions are some of the important features of this protocol.

Now days, the researchers focused one of the most fundamental heterocycles, Tetrahydropyridines (THPDs) have been the topic of passionate research for their stupendous biological importance. A quite large number of methods have been developed by the very big efforts of chemists to get libraries containing important THPDs and their derivatives in an efficient, ecological friendly and efficient one pot Multi Component Reaction (MCR). On the source of the reported text till now, tetrahydropyridines (THPDs) have established serious attention as promising structure for several natural and synthetic molecules and have been known as the primary core structure of large diversity of biologically important mainly nitrogen and phosphorus containing heterocyclic compounds. Among the various methods reported for the synthesis of THPDs derivatives viz [1]. Proline mediated cascade Mannich-type cyclization [2], palladium-catalysed cyclizations [3], annulations of aziridines, cyclization of N-allyl amino-substituted adducts [4] and one-pot multicomponent reactions (MCRs) [5].



Synthesis of tetrahydropyridine phosphonate derivatives

## References

1. Lebold T. P., Leduc A. B., Kerr M. A. Zn(II)-Catalyzed Synthesis of Piperidines from Propargyl Amines and Cyclopropanes // *Org. Lett.* American Chemical Society. 2009. Vol. 11, № 16. P. 3770–3772.
2. Proline-Mediated Enantioselective Construction of Tetrahydropyridines *via* a Cascade Mannich Type / Intramolecular Cyclization Reaction / R. Han [et al.] // *Adv. Synth. Catal.* John Wiley & Sons, Ltd. 2008. Vol. 350, № 10. P. 1474–1478.
3. Yoshida M., Kinoshita K., Namba K. Synthesis of functionalized 2-vinyl-2,3-dihydropyrroles and 3-methylene-1,2,3,4-tetrahydropyridines by palladium-catalyzed cyclization of  $\beta$ -enaminocarbonyl compounds with allylic bisacetates // *Org. Biomol. Chem.* Royal Society of Chemistry. 2014. Vol. 12, № 15. P. 2394–2403.

4. Distinct reactivity of Morita – Baylis – Hillman acetates as a novel C2 component in amine-catalyzed [2 + 2 + 2] and [2 + 4] annulations / R. Chen [et al.] // Chem. Commun. Royal Society of Chemistry. 2013. Vol. 49, № 34. P. 3543.

5. Organocatalytic asymmetric multicomponent reactions of aromatic aldehydes and anilines with  $\beta$ -ketoesters: facile and atom-economical access to chiral tetrahydropyridines / X. Li [et al.] // Chem. Commun. Royal Society of Chemistry. 2013. Vol. 49, № 14. P. 1401.